EPJ Web of Conferences will be set by the publisher
DOI: will be set by the publisher
© Owned by the authors, published by EDP Sciences, 2016

arXiv:1609.04651v1 [hep-ph] 15 Sep 2016

Strong Couplings of Three Mesons with Charm(ing) Involvement

Wolfgang Lucha[1,a], Dmitri Melikhov[1,2,3,b], Hagop Sazdjian[4,c], and Silvano Simula[5,d]

[1] *Institute for High Energy Physics, Austrian Academy of Sciences, Nikolsdorfergasse 18, A-1050 Vienna, Austria*
[2] *D. V. Skobeltsyn Institute of Nuclear Physics, M. V. Lomonosov Moscow State University, 119991, Moscow, Russia*
[3] *Faculty of Physics, University of Vienna, Boltzmanngasse 5, A-1090 Vienna, Austria*
[4] *IPN, CNRS/IN2P3, Université Paris-Sud 11, F-91406 Orsay, France*
[5] *INFN, Sezione di Roma Tre, Via della Vasca Navale 84, I-00146 Roma, Italy*

Abstract. We determine the strong couplings of three mesons that involve, at least, one η_c or J/ψ meson, within the framework of a constituent-quark model by means of relativistic dispersion formulations. For strong couplings of J/ψ mesons to two charmed mesons, our approach leads to predictions roughly twice as large as those arising from QCD sum rules.

1 Three-meson strong coupling from meson–meson transition amplitudes

We determine the strong couplings of three mesons at least one of which is one of the charmonia η_c and J/ψ, generically called $g_{PP'V}$ and $g_{PV'V}$ for pseudoscalar mesons P of mass M_P and vector mesons V of mass M_V and polarization vector ε_μ and defined, for momentum transfer $q \equiv p_1 - p_2$, by the amplitudes

$$\langle P'(p_2)\, V(q)|P(p_1)\rangle = -\frac{g_{PP'V}}{2}\,(p_1+p_2)^\mu\,\varepsilon^*_\mu(q)\,,$$
$$\langle V'(p_2)\, V(q)|P(p_1)\rangle = -g_{PV'V}\,\epsilon_{\mu\nu\rho\sigma}\,\varepsilon^{*\mu}(q)\,\varepsilon^{*\nu}(p_2)\,p_1^\rho\,p_2^\sigma\,,$$

from the residues of *poles* situated at the masses M_{P_R} and M_{V_R} of (appropriate) pseudoscalar and vector resonances P_R and V_R and contributing to *transition form factors* $F_+^{P\to P'}(q^2)$, $V^{P\to V}(q^2)$ and $A_0^{P\to V}(q^2)$, in terms of vector quark currents $j_\mu \equiv \bar q_1\,\gamma_\mu\,q_2$ and axial-vector quark currents $j_\mu^5 \equiv \bar q_1\,\gamma_\mu\,\gamma_5\,q_2$ defined by

$$\langle P'(p_2)|j_\mu|P(p_1)\rangle = F_+^{P\to P'}(q^2)\,(p_1+p_2)_\mu + \cdots\,, \qquad F_+^{P\to P'}(q^2)\Big|_{\text{pole}} = \frac{g_{PP'V_R}\,f_{V_R}}{2\,M_{V_R}\left(1-q^2/M_{V_R}^2\right)}\,,$$

$$\langle V(p_2)|j_\mu|P(p_1)\rangle = \frac{2\,V^{P\to V}(q^2)}{M_P+M_V}\,\epsilon_{\mu\nu\rho\sigma}\,\varepsilon^{*\nu}(p_2)\,p_1^\rho\,p_2^\sigma\,, \qquad V^{P\to V}(q^2)\Big|_{\text{pole}} = \frac{(M_V+M_P)\,g_{PVV_R}\,f_{V_R}}{2\,M_{V_R}\left(1-q^2/M_{V_R}^2\right)}\,,$$

$$\langle V(p_2)|j_\mu^5|P(p_1)\rangle = \mathrm{i}\,q_\mu\,(\varepsilon^*(p_2)\,p_1)\,\frac{2\,M_V}{q^2}\,A_0^{P\to V}(q^2) + \cdots\,, \qquad A_0^{P\to V}(q^2)\Big|_{\text{pole}} = \frac{g_{PP_RV}\,f_{P_R}}{2\,M_V\left(1-q^2/M_{P_R}^2\right)}\,,$$

[a] e-mail: Wolfgang.Lucha@oeaw.ac.at
[b] e-mail: dmitri_melikhov@gmx.de
[c] e-mail: sazdjian@ipno.in2p3.fr
[d] e-mail: simula@roma3.infn.it

where the P or V decay constants $f_{P,V}$ parametrize the matrix elements of the interpolating currents $j_\mu^{(5)}$

$$\langle 0|j_\mu^5|P(q)\rangle = \mathrm{i}\, f_P\, q_\mu\ , \qquad \langle 0|j_\mu|V(q)\rangle = f_V\, M_V\, \varepsilon_\mu(q)\ .$$

Such strong-coupling results may prove to be useful for studies of long-distance QCD effects in hadron decays involving charmed mesons or charmonia in the final state of a kind similar to the one in Ref. [1].

2 Quark-model-underpinned dispersion analysis of transition form factors

We describe the relevant properties of the involved strongly coupling mesons by means of a relativistic constituent-quark model [2–4]. Of course, this requires us to match the QCD currents $j_\mu^{(5)}$ to associated constituent-quark currents, which is, for heavy quarks, easily effected by introducing form factors $g_{V,A}$,

$$j_\mu = g_V\, \bar{Q}_1\, \gamma_\mu\, Q_2 + \text{other Lorentz structures}\ , \qquad j_\mu^5 = g_A\, \bar{Q}_1\, \gamma_\mu\, \gamma_5\, Q_2 + \text{other Lorentz structures}\ ,$$

for which we choose $g_V = g_A = 1$ [5] but, for light quarks, rendered rather involved [6, 7], for instance, if embedding partial axial-current conservation. For the radial meson wave functions, Gaussian shapes

$$w_{P,V}(k^2) \propto \exp\left(-\frac{k^2}{2\,\beta_{P,V}^2}\right), \qquad \int \mathrm{d}k\, k^2\, w_{P,V}^2(k^2) = 1\ ,$$

with slopes $\beta_{P,V}$ given, together with all relevant mesonic features, in Table 1 [8–13], turn out to suffice for our purposes. Table 2 lists the numerical values adopted for the masses of the constituent quarks Q. Within the framework of a relativistic dispersion formalism (reviewed, e.g., in Ref. [14]), we represent each transition form factor $\mathcal{F}(q^2) = F_+^{P\to P'}(q^2), V^{P\to V}(q^2), A_0^{P\to V}(q^2)$ by a double dispersion integral of a double spectral density $\Delta_{\mathcal{F}}(s_1, s_2, q^2)$ the one-loop contributions to which derive from Feynman graphs like the ones in Fig. 1 and each decay constant $f_{P,V}$ by a dispersion integral of a spectral density $\rho_{P,V}(s)$,

$$\mathcal{F}(q^2) = \int \mathrm{d}s_1\, \mathrm{d}s_2\, \phi_1(s_1)\, \phi_2(s_2)\, \Delta_{\mathcal{F}}(s_1, s_2, q^2)\ , \qquad f_{P,V} = \int\limits_{(m_1+m)^2}^{\infty} \mathrm{d}s\, \phi_{P,V}(s)\, \rho_{P,V}(s)\ ,$$

Table 1. Relevant parameters of the mesons: numerical values of mass M, leptonic decay constant f and slope β.

Meson	M (GeV)	f (MeV)	β (GeV)
D	1.87	206 ± 8	0.475
D^*	2.010	260 ± 10	0.48
D_s	1.97	248 ± 2.5	0.545
D_s^*	2.11	311 ± 9	0.54
η_c	2.980	394.7 ± 2.4	0.77
J/ψ	3.097	405 ± 7	0.68

Table 2. Constituent mass of each quark flavour $Q = u, d, s, c$ [5] involved in charm(ing) three-meson couplings.

Quark flavour	Quark mass m (GeV)
u	0.23
d	0.23
s	0.35
c	1.45



Figure 1. Feynman graphs of transitions among the mesons η_c, J/ψ, D, D^*, induced by the currents $\bar{c}\,\gamma_\mu\,c$ or $\bar{c}\,\gamma_\mu\,d$.

and of the wave functions of all mesons entering the corresponding one- or two-meson matrix elements

$$\phi_{P,V}(s) = \frac{\pi}{s^{3/4}} \sqrt{\frac{s^2 - (m_1^2 - m^2)^2}{2\,[s - (m_1 - m)^2]}}\; w_{P,V}\!\left(\frac{(s - m_1^2 - m^2)^2 - 4\,m_1^2\,m^2}{4\,s}\right).$$

3 Three-meson strong coupling: determination from transition amplitudes

We fix the slopes $\beta_{P,V}$ such that the decay constants $f_{P,V}$ are reproduced by their spectral representation. Equipped with these $\beta_{P,V}$ values, we deduce all strong couplings from the spectral representation of the relevant form factors $\mathcal{F}(q^2)$ derived sufficiently off the resonances at M_R ($R = P_R, V_R$), by interpolating pointwise given momentum dependences of $\mathcal{F}(q^2)$ by three-parameter ($\sigma_{1,2}, \mathcal{F}(0)$) ansätze of the form

$$\mathcal{F}(q^2) = \frac{\mathcal{F}(0)}{1 - \sigma_1\,q^2/M_R^2 + \sigma_2\,q^4/M_R^4}\;\frac{1}{1 - q^2/M_R^2}\,, \qquad \mathrm{Res}\,\mathcal{F}(M_R^2) = \frac{\mathcal{F}(0)}{1 - \sigma_1 + \sigma_2}\,, \qquad R = P_R, V_R\,,$$

and extrapolating $\mathcal{F}(q^2)$ to the poles at $q^2 = M_R^2$, where the strong couplings emerge from the residues. Using $\sigma_{1,2}$, $\mathcal{F}(0)$, and M_R as fit parameters, all arising masses M_R come close to the known resonances. Quite generally, a given strong coupling may show up in and therefore can be extracted from more than one meson–meson transition form factor, for example, $g_{\eta_c\eta_c\psi}$ from $F_+^{\eta_c>\eta_c}$ or $A_0^{\eta_c>\psi}$ (see Fig. 2) [15–17], $g_{DD\psi}$ from $F_+^{D>D}$ or $A_0^{D>\psi}$ (see Fig. 3(a)) [15–17] and $g_{DD^*\eta_c}$ from $F_+^{\eta_c>D}$, $A_0^{\eta_c>D^*}$ or $A_0^{D>D^*}$ (see Fig. 3(b)) [15–17]; for further examples of such multiple involvements, consult Tables III, V, and VI of Ref. [15].



Figure 2. Behaviour of the off-shell η_c-η_c-J/ψ strong coupling $g_{\eta_c\eta_c\psi}$ with increasing resonance-mass-normalized momentum transfer $x \equiv q^2/M_R^2$ for the transition of η_c to the η_c (solid red line) or the J/ψ meson (dotted blue line).



Figure 3. Behaviour of the "off-shell" D-D-J/ψ and D-D^*-η_c strong couplings $g_{DD\psi}$ and $g_{DD^*\eta_c}$, respectively, with increasing "resonance-mass-normalized" momentum transfer $x \equiv q^2/M_R^2$: (a) $g_{D\hat{D}\psi}(x) = 2\,M_\psi\,(1-x)\,A_0^{D>\psi}(q^2)/\,f_D$ (blue dotted line and squares ■) and $g_{DD\hat{\psi}}(x) = 2\,M_\psi\,(1-x)\,F_+^{D>D}(q^2)/\,f_\psi$ (red solid line and triangles ▲), relying on interpolation (blue or red lines) or not (squares ■, triangles ▲); (b) $g_{D\hat{D}^*\eta_c}(x)$ (solid red line), $g_{DD^*\hat{\eta}_c}(x)$ (dotted blue line) and $g_{\hat{D}D^*\eta_c}(x)$ (dashed green line). For each transition, the relevant resonance, R, is identified by a circumflex.

4 Strong coupling predictions from relativistic constituent-quark approach

We collect our emerging strong-coupling findings — extracted, in the case of multipresence of one and the same three-meson coupling in more than one meson–meson transition amplitude, by a combined fit — in Table 3: Strange quark content instead of a down quark implies a reduction of the involved strong couplings, by roughly 10%. Confronting, in Table 4, our $D_{(s)}$-$D_{(s)}^{(*)}$-J/ψ predictions with QCD sum-rule outcomes [18–20], the QCD sum-rule estimates prove to be lower than ours [15–17] by a factor of two.

Table 3. Charm(ing) three-meson strong couplings: quark-model-based dispersion-approach outcomes [15–17].

$PP'V$ Coupling	Strong coupling $g_{PP'V}$	$PV'V$ coupling	Strong coupling $g_{PV'V}$ (GeV^{-1})
η_c-η_c-J/ψ	25.8 ± 1.7	η_c-J/ψ-J/ψ	10.6 ± 1.5
D-D-J/ψ	26.04 ± 1.43	D-D^*-J/ψ	10.7 ± 0.4
D-D^*-η_c	15.51 ± 0.45	D^*-D^*-η_c	9.76 ± 0.32
D_s-D_s-J/ψ	23.83 ± 0.78	D_s-D_s^*-J/ψ	9.6 ± 0.8
D_s-D_s^*-η_c	14.15 ± 0.52	D_s^*-D_s^*-η_c	8.27 ± 0.37

Table 4. Strong couplings of the J/ψ meson to two charmed mesons: relativistic quark model vs. QCD sum rule.

Coupling \ Approach	Quark model [15–17]	QCD sum rules	References
D-D-J/ψ	26.04 ± 1.43	11.6 ± 1.8	[18]
D-D^*-J/ψ	(10.7 ± 0.4) GeV^{-1}	(4.0 ± 0.6) GeV^{-1}	[18]
D_s-D_s-J/ψ	23.83 ± 0.78	$11.96^{+1.34}_{-1.16}$	[19]
D_s-D_s^*-J/ψ	(9.6 ± 0.8) GeV^{-1}	$(4.30^{+1.53}_{-1.22})$ GeV^{-1}	[20]

Acknowledgements

D. M. would like to express gratitude for support by the Austrian Science Fund (FWF) under project P29028-N27.

References

[1] P. Santorelli, Phys. Rev. D **77** (2008) 074012, arXiv:hep-ph/0703232.
[2] W. Lucha, F. F. Schöberl, and D. Gromes, Phys. Rep. **200** (1991) 127.
[3] F. Cardarelli, E. Pace, G. Salmè, and S. Simula, Phys. Lett. B **357** (1995) 267, arXiv:nucl-th/ 9507037.
[4] R. N. Faustov and V. O. Galkin, Z. Phys. C **66** (1995) 119.
[5] D. Melikhov and B. Stech, Phys. Rev. D **62** (2000) 014006, arXiv:hep-ph/0001113.
[6] D. Melikhov and B. Stech, Phys. Rev. D **74** (2006) 034022, arXiv:hep-ph/0606203.
[7] W. Lucha, D. Melikhov, and S. Simula, Phys. Rev. D **74** (2006) 054004, arXiv:hep-ph/0606281.
[8] C. T. H. Davies *et al.*, Phys. Rev. D **82** (2010) 114504, arXiv:1008.4018 [hep-lat].
[9] W. Lucha, D. Melikhov, and S. Simula, Phys. Lett. B **701** (2011) 82, arXiv:1101.5986 [hep-ph].
[10] D. Bečirević *et al.*, J. High Energy Phys. **1202** (2012) 042, arXiv:1201.4039 [hep-lat].
[11] G. C. Donald *et al.* (HPQCD Coll.), Phys. Rev. D **86** (2012) 094501, arXiv:1208.2855 [hep-lat].
[12] W. Lucha, D. Melikhov, and S. Simula, Phys. Lett. B **735** (2014) 12, arXiv:1404.0293 [hep-ph].
[13] K. A. Olive *et al.* (Particle Data Group), Chin. Phys. C **38** (2014) 090001 and 2015 update.
[14] D. Melikhov, Eur. Phys. J. direct **C4** (2002) 2, arXiv:hep-ph/0110087.
[15] W. Lucha, D. Melikhov, H. Sazdjian, and S. Simula, Phys. Rev. D **93** (2016) 016004, **93** (2016) 019902(E), arXiv:1506.09213 [hep-ph].
[16] W. Lucha, D. Melikhov, H. Sazdjian, and S. Simula, Proc. Sci., EPS-HEP 2015 (2015) 428, arXiv:1509.03089 [hep-ph].
[17] W. Lucha, D. Melikhov, H. Sazdjian, and S. Simula, preprint HEPHY-PUB 970/16 (2016), arXiv:1607.05569 [hep-ph].
[18] R. D. Matheus *et al.*, Int. J. Mod. Phys. E **14** (2005) 555.
[19] B. Osório Rodrigues, M. E. Bracco, and M. Chiapparini, Nucl. Phys. A **929** (2014) 143, arXiv: 1309.1637 [hep-ph].
[20] B. Osório Rodrigues *et al.*, Eur. Phys. J. A **51** (2015) 28, arXiv:1501.03088 [hep-ph].